### §520.903c

## §520.903c [Reserved]

## § 520.903d Febantel-praziquantel paste.

- (a) *Specifications*. Each gram of paste contains 34 milligrams of febantel and 3.4 milligrams of praziquantel.
- (b) *Sponsor*. See No. 000859 in §510.600(c) of this chapter.
- (c) Conditions of use—(1) Amount—(i) Dogs and cats (over 6 months of age): 10 milligrams of febantel and 1 milligram of praziquantel per kilogram of body weight (1 gram of paste per 7.5 pounds body weight) administered by mouth or in the food once daily for 3 days.
- (ii) Puppies and kittens (less than 6 months of age): 15 milligrams of febantel and 1.5 milligrams of praziquantel per kilogram of body weight (1 gram of paste per 5 pounds body weight) administered by mouth on a full stomach once daily for 3 days.
- (2) Indications for use. (i) Dogs and puppies: For removal of hookworms (Ancylostoma caninum and Uncinaria stenocephala), whipworms (Trichuris vulpis), ascarids (Toxocara canis and Toxascaris leonina), and tapeworms (Dipylidium caninum and Taenia pisiformis).
- (ii) Cats and kittens: For removal of hookworms (Ancylostoma tubaeforme), ascarids (Toxocara cati) and tapeworms (Dipylidium caninum and Taenia taeniaeformis).
- (3) *Limitations*. Do not use in pregnant animals. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (4) Special considerations. Consider alternative therapy or use with caution in animals with pre-existing liver or kidney dysfunction.
- [50 FR 19167, May 7, 1985, as amended at 53 FR 48533, Dec. 1, 1988; 56 FR 50813, Oct. 9, 1991]

## § 520.903e Febantel tablets.

- (a) Specifications. Each scored tablet contains 27.2 milligrams of febantel for use in dogs, puppies, cats, and kittens or 163.3 milligrams of febantel for use in dogs, puppies, and cats.
- (b) Sponsor. See 000859 in 510.600(c)(2) of this chapter.
- (c) Conditions of use—(1) Amount—(i) Dogs and cats. Ten milligrams per kilo-

gram body weight. Administer once daily for 3 consecutive days.

- (ii) Puppies and kittens fewer than 6 months of age. Fifteen milligrams per kilogram body weight. Administer once daily for 3 consecutive days.
- (2) Indications for use. (i) For removal of hookworms (Ancylostoma caninum and Uncinaria stenocephala), ascarids (Toxocara canis and Toxascaris leonina) and whipworms (Trichuris vulpis) in dogs and puppies.
- (ii) For removal of hookworms (Ancylostoma tubaeforme) and ascarids (Toxocara cati) in cats and kittens.
- (3) Limitations. Do not use in pregnant animals. Consider alternative therapy or use with caution in animals with preexisting liver or kidney dysfunction. Administer to puppies and kittens on a full stomach. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[56 FR 50655, Oct. 8, 1991]

# § 520.905 Fenbendazole oral dosage forms.

#### §520.905a Fenbendazole suspension.

- (a) Specifications. Each milliliter of suspension contains 100 milligrams (mg) fenbendazole.
- (b) Sponsor. See No. 000061 in §510.600(c) of this chapter.
- (c) Related tolerances. See §556.275 of this chapter.
- (d) Special considerations. (1) See §500.25 of this chapter.
- (2) Fenbendazole suspension 10 percent and approved forms of trichlorfon, when used concomitantly for treating the indications provided in paragraph (e) of this section and for treating infections of stomach bot as provided in §520.2520, have been shown to be compatible and not to interfere with one another.
- (e) Conditions of use—(1) Horses—(i) Amount. Administer orally 5 mg per kilogram (/kg) (2.3 mg per pound (/lb)) for the control of large strongyles, small strongyles, and pinworms; 10 mg/kg for the control of ascarids.
- (ii) Indications for use. For the control of large strongyles (Strongylus edentatus, S. equinus, S. vulgaris), small strongyles (Cyanthostomum spp., Cylicocyclus spp., Cylicostephanus spp., Triodontophorus spp.), pinworms